

Specimen Number	Patient ID	Control Number	Account Number	Account Phone Number	Route
Patient Last Name			Account Address		
Patient First Name		Patient Middle Name			
Patient SS#	Patient Phone	Total Volume			
Age (Y/M/D)	Date of Birth	Sex	Fasting		
Patient Address			Additional Information		
Date and Time Collected	Date Entered	Date and Time Reported	Physician Name	NPI	Physician ID

Tests Ordered					
ANCA Panel					

TESTS	RESULT	FLAG	UNITS	REFERENCE INTERVAL	LAB
-------	--------	------	-------	--------------------	-----

ANCA Panel

Antimyeloperoxidase (MPO) Abs	<9.0		U/mL	0.0 - 9.0	01
Antiproteinase 3 (PR-3) Abs	<3.5		U/mL	0.0 - 3.5	01
Cytoplasmic (C-ANCA)	<1:20		titer	Neg:<1:20	01
Perinuclear (P-ANCA)	<1:20		titer	Neg:<1:20	01

The presence of positive fluorescence exhibiting P-ANCA or C-ANCA patterns alone is not specific for the diagnosis of Wegener's Granulomatosis (WG) or microscopic polyangiitis. Decisions about treatment should not be based solely on ANCA IFA results. The International ANCA Group Consensus recommends follow up testing of positive sera with both PR-3 and MPO-ANCA enzyme immunoassays. As many as 5% serum samples are positive only by EIA.
Ref. AM J Clin Pathol 1999;111:507-513.

Atypical pANCA	<1:20		titer	Neg:<1:20	01
The atypical pANCA pattern has been observed in a significant percentage of patients with ulcerative colitis, primary sclerosing cholangitis and autoimmune hepatitis.					

--

--	--	--

DUPLICATE FINAL REPORT

Page 1 of 1