DOB:

## **Patient Report**

Ordering Physician:

labcorp

Patient ID:

Specimen ID: Sex:

#### Ordered Items: Actin (Smooth Muscle) Antibody; Liver-Kidney Microsomal Ab; Antinuclear Antibodies, IFA; Venipuncture

Date Collected: Date Received: Date Reported: Fasting:

### **Actin (Smooth Muscle) Antibody**

Test	<b>Current Result and Flag</b>	Previous Result and Date	Units	Reference Interval		
Actin (Smooth Muscle) Antibody 01	5		Units	0-19		
		Negative	0 - 19			
		Weak positive	20 - 30			
		Moderate to strong posi	tive >30			
	Actin Antil	oodies are found in 52-85% of	patients with			
	autoimmune hepatitis or chronic active hepatitis and					
	in 22% of patients with primary biliary cirrhosis.					

# **Liver-Kidney Microsomal Ab**

Test	<b>Current Result and Flag</b>	Previous Result and Date	Units	Reference Interval
Liver-Kidney Microsomal Ab 01	1.0		Units	0.0-20.0
		Negative	0.0 - 20.0	
		Equivocal	20.1 - 24.9	
		Positive	>24.9	
	LKM type 1 an	tibodies are detected in pati	ients with	
	autoimmune he	patitis type 2 and in up to 8	3% of	
	patients with	chronic HCV infection.		

# **Antinuclear Antibodies, IFA**

Test	Current Result and Flag	Previous Result and Date	Units	Reference Interval
Antinuclear Antibodies, IFA 02	Negative			
		Negat	ive <1:80	
		Borde	rline 1:80	
		Posit	ive >1:80	
	ICAP nomenclature: AC-0			
	For more information about H	ep-2 cell patterns use		
	ANApatterns.org, the officia	official website for the International lear Antibody (ANA) Patterns (ICAP).		
	Consensus on Antinuclear Anti			

#### Disclaime

The Previous Result is listed for the most recent test performed by Labcorp in the past 3 years where there is sufficient patient demographic data to match the result to the patient.

#### Icon Legend

labcorp Final Report Page 1 of 1