

Patient ID: DOB:
 Specimen ID: Age:
 Sex:

Patient Report

Ordering Physician:



Ordered Items: **CBC With Differential/Platelet; CEA; Prostate-Specific Ag; Venipuncture**

Date Collected: Date Received: Date Reported: Fasting:

CBC With Differential/Platelet

| Test | Current Result and Flag | Previous Result and Date | Units | Reference Interval |
|--------------------------------------|-------------------------|--------------------------|----------|--------------------|
| WBC ⁰¹ | 4.1 | | x10E3/uL | 3.4-10.8 |
| RBC ⁰¹ | 4.39 | | x10E6/uL | 4.14-5.80 |
| Hemoglobin ⁰¹ | 11.9 | | g/dL | 13.0-17.7 |
| Hematocrit ⁰¹ | 37.4 | | % | 37.5-51.0 |
| MCV ⁰¹ | 85 | | fL | 79-97 |
| MCH ⁰¹ | 27.1 | | pg | 26.6-33.0 |
| MCHC ⁰¹ | 31.8 | | g/dL | 31.5-35.7 |
| RDW ⁰¹ | 13.3 | | % | 11.6-15.4 |
| Platelets ⁰¹ | 271 | | x10E3/uL | 150-450 |
| Neutrophils ⁰¹ | 46 | | % | Not Estab. |
| Lymphs ⁰¹ | 42 | | % | Not Estab. |
| Monocytes ⁰¹ | 8 | | % | Not Estab. |
| Eos ⁰¹ | 3 | | % | Not Estab. |
| Basos ⁰¹ | 1 | | % | Not Estab. |
| Neutrophils (Absolute) ⁰¹ | 1.9 | | x10E3/uL | 1.4-7.0 |
| Lymphs (Absolute) ⁰¹ | 1.7 | | x10E3/uL | 0.7-3.1 |
| Monocytes(Absolute) ⁰¹ | 0.3 | | x10E3/uL | 0.1-0.9 |
| Eos (Absolute) ⁰¹ | 0.1 | | x10E3/uL | 0.0-0.4 |
| Baso (Absolute) ⁰¹ | 0.0 | | x10E3/uL | 0.0-0.2 |
| Immature Granulocytes ⁰¹ | 0 | | % | Not Estab. |
| Immature Grans (Abs) ⁰¹ | 0.0 | | x10E3/uL | 0.0-0.1 |

CEA

| Test | Current Result and Flag | Previous Result and Date | Units | Reference Interval |
|-------------------|-------------------------|--|-------|--------------------|
| CEA ⁰¹ | 3.0 | | ng/mL | 0.0-4.7 |
| | | Nonsmokers | <3.9 | |
| | | Smokers | <5.6 | |
| | | Roche Diagnostics Electrochemiluminescence Immunoassay (ECLIA) | | |
| | | Values obtained with different assay methods or kits cannot be used interchangeably. Results cannot be interpreted as absolute evidence of the presence or absence of malignant disease. | | |

Prostate-Specific Ag

| Test | Current Result and Flag | Previous Result and Date | Units | Reference Interval |
|------------------------------------|-------------------------|---|-------|--------------------|
| Prostate Specific Ag ⁰¹ | 1.5 | | ng/mL | 0.0-4.0 |
| | | Roche ECLIA methodology. | | |
| | | According to the American Urological Association, Serum PSA should decrease and remain at undetectable levels after radical | | |

Patient ID:
Specimen ID:

DOB:
Age:
Sex:

Patient Report

Ordering Physician:



Prostate-Specific Ag (Cont.)

prostatectomy. The AUA defines biochemical recurrence as an initial PSA value 0.2 ng/mL or greater followed by a subsequent confirmatory PSA value 0.2 ng/mL or greater.

Values obtained with different assay methods or kits cannot be used interchangeably. Results cannot be interpreted as absolute evidence of the presence or absence of malignant disease.

Disclaimer

The Previous Result is listed for the most recent test performed by Labcorp in the past 5 years where there is sufficient patient demographic data to match the result to the patient. Results from certain tests are excluded from the Previous Result display.

Icon Legend

▲ Out of Reference Range ■ Critical or Alert

Performing Labs

Patient Details

Phone:
Date of Birth:
Age:
Sex:
Patient ID:
Alternate Patient ID:

Physician Details

Request A Test
7027 Mill Road Suite 201, BRECKSVILLE, OH,
44141

Phone: **888-732-2348**
Physician ID:
NPI:

Specimen Details

Specimen ID:
Control ID:
Alternate Control Number:
Date Collected:
Date Received:
Date Entered:
Date Reported:
Rte: