

Specimen Number	Patient ID	Control Number	Account Number	Account Phone Number	Route
Patient Last Name		Account Address			
Patient First Name		Patient Middle Name			
Patient SS#	Patient Phone	Total Volume			
Age (Y/M/D)	Date of Birth	Sex	Fasting		
Patient Address			Additional Information		
Date and Time Collected	Date Entered	Date and Time Reported	Physician Name	NPI	Physician ID

Tests Ordered

Celiac Disease Complete Panel; Venipuncture

TESTS	RESULT	FLAG	UNITS	REFERENCE INTERVAL	LAB
-------	--------	------	-------	--------------------	-----

Celiac Disease Complete Panel

Immunoglobulin A, Qn, Serum	157		mg/dL	44 - 189	01
Deamidated Gliadin Abs, IgA	5		units	0 - 19	02
		Negative		0 - 19	
		Weak Positive		20 - 30	
		Moderate to Strong Positive		>30	
Deamidated Gliadin Abs, IgG	3		units	0 - 19	02
		Negative		0 - 19	
		Weak Positive		20 - 30	
		Moderate to Strong Positive		>30	
t-Transglutaminase (tTG) IgA	<2		U/mL	0 - 3	02
		Negative		0 - 3	
		Weak Positive		4 - 10	
		Positive		>10	

Tissue Transglutaminase (tTG) has been identified as the endomysial antigen. Studies have demonstrated that endomysial IgA antibodies have over 99% specificity for gluten sensitive enteropathy.

t-Transglutaminase (tTG) IgG	2		U/mL	0 - 5	02
		Negative		0 - 5	
		Weak Positive		6 - 9	
		Positive		>9	

--	--	--