

Patient Information	Specimen Information	Client Information
<b>DOB:</b> <b>AGE:</b> Gender: Phone: Patient ID:	Specimen: Requisition: Lab Ref #:  Collected: Received: Reported:	

**COMMENTS:**            FASTING: YES

Test Name	In Range	Out Of Range	Reference Range	Lab
GLUCOSE	98		65-99 mg/dL	QBA

Fasting reference interval

HEMOGLOBIN A1c WITH eAG				QBA
HEMOGLOBIN A1c	5.4		<5.7 % of total Hgb	

According to ADA guidelines, hemoglobin A1c <7.0% represents optimal control in non-pregnant diabetic patients. Different metrics may apply to specific patient populations. Standards of Medical Care in Diabetes-2013. Diabetes Care. 2013;36:s11-s66

For the purpose of screening for the presence of diabetes

<5.7%	Consistent with the absence of diabetes
5.7-6.4%	Consistent with increased risk for diabetes (prediabetes)
>or=6.5%	Consistent with diabetes

This assay result is consistent with a decreased risk of diabetes.

Currently, no consensus exists for use of hemoglobin A1c for diagnosis of diabetes for children.

eAG (mg/dL)	108	(calc)
eAG (mmol/L)	6.0	(calc)

**PERFORMING SITE:**

QBA QUEST DIAGNOSTICS-BALTIMORE, 1901 SULPHUR SPRING ROAD, BALTIMORE, MD 21227-2943 Laboratory Director: EDGAR G KHALLUF, MD, CLIA: 21D0218877