**Patient Report** 

Ordering Physician:

DOB: Age: Sex:



Ordered Items: G6PD, Qn, Bld and Red Cell Count; Venipuncture

Date Collected: Date Received: Date Reported: Fasting:

## G6PD,Qn,Bld and Red Cell Count

Patient ID: Specimen ID:

Test	Current Result and Flag	<b>Previous Result and Date</b>	Units	Reference Interval
RBC 02	4.27		x10E6/uL	3.77-5.28
G-6-PD, Quant	245		U/10E12 RBC	127-427
Comment: 01				

When decreased, G-6-PD, Quant. values are associated with acute hemolytic anemia when deficient individuals are exposed to oxidative stress, such as with certain medications (e.g., primaquine), infection, or ingestion of fava beans. Caution: In patients with acute hemolysis (e.g., abnormally low RBC values), testing for G-6-PD may be falsely normal because older

erythrocytes with a higher enzyme deficiency have been hemolyzed. Young erythrocytes and reticulocytes have normal or near-normal enzyme activity. Normal values of G-6-PD may be measured for several weeks following a hemolytic event.

## Disclaimer

The Previous Result is listed for the most recent test performed by Labcorp in the past 3 years where there is sufficient patient demographic data to match the result to the patient.

## Icon Legend

Phone:

Sex:

ID:

Date of Birth: Age:

Patient ID: Alternate Patient

**Patient Details** Physician Details Specimen Details

> Specimen ID: Control ID:

Phone: Alternate Control Number: Physician ID: Date Collected: NPI:

Date Received: Date Entered:

Date Reported:

Rte:

labcorp Final Report Page 1 of 1