



Report Status: Final NOMIKOS, JOHN

Patient Information	Specimen Information	Client Information
DOB: AGE: Gender: Phone: Patient ID: Health ID:	Specimen: Requisition: Lab Ref #: Collected: Received: Reported:	REQUEST A TEST - PWN HEAL TH 7027 MILL RD STE 201 BRECKSVILLE, OH 44141-1852

Test Name	In Range	Out Of Range	Reference Range	Lab
HSV 1/2 IGG, W/REFL	119-124 119-134-134	Constitution Scott Institution	Miles to the common the control of t	CB
HSV2 INHIBITION				
HSV 1 IGG, TYPE SPECIFIC				
AB	<0.90		index	
HSV 2 IGG, TYPE SPECIFIC				
AB	<0.90		index	
	Index	Interpretation		

	<0.90	Negative		
	0.90-1.09	Equivocal		
	>1.09	Positive		

This assay utilizes recombinant type-specific antigens to differentiate HSV-1 from HSV-2 infections. A positive result cannot distinguish between recent and past infection. If recent HSV infection is suspected but the results are negative or equivocal, the assay should be repeated in 4-6 weeks. The performance characteristics of the assay have not been established for pediatric populations, immunocompromised patients, or neonatal screening.

See Endnote 1

Endnote 1

For additional information, please refer to http://education.QuestDiagnostics.com/faq/FAQ118 (This link is being provided for informational/educational purposes only.)

PERFORMING SITE:

CB QUEST DIAGNOSTICS WOOD DALE, 1355 MITTEL BOULEVARD, WOOD DALE, IL 60191-1024 Laboratory Director: ANTHONY V. THOMAS, MD, CLIA: 14D0417052