



Patient Information	Specimen Information	Client Information
DOB: AGE: Gender: Phone: Patient ID: Health ID:	Specimen: Requisition: Lab Ref #: Collected: Received: Reported:	

Test Name	In Range	Out Of Range	Reference Range	Lab
HSV 1/2 IGG, TYPE SPECIFIC AB				QHO
HSV 1 IGG, TYPE SPECIFIC AB	<0.90		index	
HSV 2 IGG, TYPE SPECIFIC AB	<0.90		index	
	Index	Interpretation		
	<0.90	Negative		
	0.90-1.09	Equivocal		
	>1.09	Positive		

This assay utilizes recombinant type-specific antigens to differentiate HSV-1 from HSV-2 infections. A positive result cannot distinguish between recent and past infection. If recent HSV infection is suspected but the results are negative or equivocal, the assay should be repeated in 4-6 weeks. The performance characteristics of the assay have not been established for pediatric populations, immunocompromised patients, or neonatal screening.

HSV 1/2 AB (IGM), IFA				QHO
W/REL TO TITER				
HSV 1 IGM SCREEN	NEGATIVE		NEGATIVE	
HSV 2 IGM SCREEN	NEGATIVE		NEGATIVE	

The IFA procedure for measuring IgM antibodies to HSV 1 and HSV 2 detects both type-common and type-specific HSV antibodies. Thus, IgM reactivity to both HSV 1 and HSV 2 may represent cross-reactive HSV antibodies rather than exposure to both HSV 1 and HSV 2. This assay was developed and its performance characteristics have been determined by Quest Diagnostics. Performance characteristics refer to the analytical performance of the test.