

Specimen Number	Patient ID	Control Number	Account Number	Account Phone Number	Route
Patient Last Name			Account Address		
Patient First Name		Patient Middle Name			
Patient SS#	Patient Phone	Total Volume			
Age (Y/M/D)	Date of Birth	Sex	Fasting		
Patient Address			Additional Information		
Date and Time Collected	Date Entered	Date and Time Reported	Physician Name	NPI	Physician ID

Tests Ordered					
Inflammatory Bowel Disease-IBD					

TESTS	RESULT	FLAG	UNITS	REFERENCE INTERVAL	LAB
-------	--------	------	-------	--------------------	-----

Inflammatory Bowel Disease-IBD					
Saccharomyces cerevisiae, IgG	<20.0		Units	0.0 - 24.9	01
			Negative	<20.0	
			Equivocal	20.1 - 24.9	
			Positive	>or= 25.0	
Saccharomyces cerevisiae, IgA	<20.0		Units	0.0 - 24.9	01
			Negative	<20.0	
			Equivocal	20.1 - 24.9	
			Positive	>or= 25.0	

IgA and IgG antibody testing for *S. cerevisiae* is useful adjunct testing for differentiating Crohn's disease and ulcerative colitis. Close to 80% of Crohn's disease patients are positive for either IgA or IgG. In ulcerative colitis, less than 15% are positive for IgG and less than 2% are positive for IgA. Fewer than 5% are positive for either IgG or IgA antibody, and no healthy controls had antibody for both.

Atypical pANCA	Negative		titer	Neg:<1:20	01
The atypical pANCA pattern has been observed in a significant percentage of patients with ulcerative colitis, primary sclerosing cholangitis and autoimmune hepatitis.					
	ASCA+/PANCA- Suggestive of Crohn's disease				
	ASCA-/PANCA+ Suggestive of Ulcerative colitis				

--

--	--	--