

Specimen ID:
 Control ID:

Phone:

Rte:



Patient Details	Specimen Details	Physician Details
DOB: Age(y/m/d): Gender: SSN: Patient ID:	Date collected: Date received: Date entered: Date reported:	Ordering: Referring: ID: NPI:

General Comments & Additional Information

 Alternate Control Number:
 Total Volume:

 Alternate Patient ID:
 Fasting:

Ordered Items

Lyme, Line Blot, Serum; Drawing Fee

TESTS	RESULT	FLAG	UNITS	REFERENCE INTERVAL	LAB
-------	--------	------	-------	--------------------	-----

Lyme, Line Blot, Serum

Lyme Ab IgG by WB:					01
IgG P93 Ab.	Absent				01
IgG P66 Ab.	Absent				01
IgG P58 Ab.	Absent				01
IgG P45 Ab.	Absent				01
IgG P41 Ab.	Absent				01
IgG P39 Ab.	Absent				01
IgG P30 Ab.	Absent				01
IgG P28 Ab.	Absent				01
IgG P23 Ab.	Absent				01
IgG P18 Ab.	Absent				01
Lyme IgG WB Interp.	Negative				01

Positive: 5 of the following
 Borrelia-specific bands:
 18, 23, 28, 30, 39, 41, 45, 58,
 66, and 93.

Negative: No bands or banding
 patterns which do not
 meet positive criteria.

Lyme Ab IgM by WB:					01
IgM P41 Ab.	Absent				01
IgM P39 Ab.	Absent				01
IgM P23 Ab.	Absent				01
Lyme IgM WB Interp.	Negative				01

Note: An equivocal or positive EIA result followed by a negative Western Blot result is considered **NEGATIVE**. An equivocal or positive EIA result followed by a positive Western Blot is considered **POSITIVE** by the CDC.

Positive: 2 of the following bands: 23, 39 or 41

Negative: No bands or banding patterns which do not meet positive criteria.

Criteria for positivity are those recommended by CDC/ASTPHLD.

FINAL REPORT

Patient:
DOB:

Patient ID:

Control ID:

Specimen ID:
Date collected:

TESTS	RESULT	FLAG	UNITS	REFERENCE INTERVAL	LAB
-------	--------	------	-------	--------------------	-----

p23=Osp C, p41=flagellin

Note:

Sera from individuals with the following may cross react in the Lyme Western Blot assays: other spirochetal diseases (periodontal disease, leptospirosis, relapsing fever, yaws, and pinta); connective autoimmune (Rheumatoid Arthritis and Systemic Lupus Erythematosus and also individuals with Antinuclear Antibody); other infections (Rocky Mountain Spotted Fever; Epstein-Barr Virus, and Cytomegalovirus).