



Patient Information	Specimen Information	Client Information
DOB: Gender: Phone: Patient ID: Health ID:	Specimen: Requisition: Lab Ref #: Collected: Received: Reported:	

COMMENTS: FASTING:

Test Name	In Range	Out Of Range	Reference Range	Lab
MYELOPEROXIDASE ANTIBODY	<1.0	Value	Interpretation	NL1
		-----	-----	
		<1.0	No Antibody Detected	
		> or = 1.0	Antibody Detected	

Autoantibodies to myeloperoxidase (MPO) are commonly associated with the following small-vessel vasculitides: microscopic polyangiitis, polyarteritis nodosa, Churg-Strauss syndrome, necrotizing and crescentic glomerulonephritis and occasionally granulomatosis with polyangiitis (GPA, Wegener's). The perinuclear IFA pattern, (p-ANCA) is based largely on autoantibody to myeloperoxidase which serves as the primary antigen. These autoantibodies are present in active disease.

PERFORMING SITE:

AMD QUEST DIAGNOSTICS/NICHOLS CHANTILLY, 14225 NEWBROOK DRIVE, CHANTILLY, VA 20151-2228 Laboratory Director: PATRICK W. MASON,MD,PHD, CLIA: 49D0221801
 NL1 QUEST DIAGNOSTICS LLC, 200 FOREST STREET 3RD FLOOR,SUITE B, MARLBOROUGH, MA 01752-3023 Laboratory Director: SALIM E KABAWAT,MD, CLIA: 22D0076229