

Ordered Items: **CBC With Differential/Platelet; Lipid Panel; Testosterone,Free and Total; Estradiol; Prostate-Specific Ag; Blood Drawing**

Date Collected:	Date Received:	Date Reported:	Fasting:
-----------------	----------------	----------------	----------

CBC With Differential/Platelet

Test	Current Result and Flag	Previous Result and Date	Units	Reference Interval
WBC ⁰¹	8.5		x10E3/uL	3.4-10.8
RBC ⁰¹	5.19		x10E6/uL	4.14-5.80
Hemoglobin ⁰¹	15.6		g/dL	13.0-17.7
Hematocrit ⁰¹	46.1		%	37.5-51.0
MCV ⁰¹	89		fL	79-97
MCH ⁰¹	30.1		pg	26.6-33.0
MCHC ⁰¹	33.8		g/dL	31.5-35.7
RDW ⁰¹	12.7		%	11.6-15.4
Platelets ⁰¹	231		x10E3/uL	150-450
Neutrophils ⁰¹	70		%	Not Estab.
Lymphs ⁰¹	20		%	Not Estab.
Monocytes ⁰¹	7		%	Not Estab.
Eos ⁰¹	2		%	Not Estab.
Basos ⁰¹	1		%	Not Estab.
Neutrophils (Absolute) ⁰¹	6.0		x10E3/uL	1.4-7.0
Lymphs (Absolute) ⁰¹	1.7		x10E3/uL	0.7-3.1
Monocytes(Absolute) ⁰¹	0.6		x10E3/uL	0.1-0.9
Eos (Absolute) ⁰¹	0.2		x10E3/uL	0.0-0.4
Baso (Absolute) ⁰¹	0.1		x10E3/uL	0.0-0.2
Immature Granulocytes ⁰¹	0		%	Not Estab.
Immature Grans (Abs) ⁰¹	0.0		x10E3/uL	0.0-0.1

Lipid Panel

Test	Current Result and Flag	Previous Result and Date	Units	Reference Interval
Cholesterol, Total ⁰¹	146		mg/dL	100-199
Triglycerides ⁰¹	146		mg/dL	0-149
HDL Cholesterol ⁰¹	45		mg/dL	>39
VLDL Cholesterol Cal	25		mg/dL	5-40
LDL Chol Calc (NIH)	76		mg/dL	0-99

Testosterone,Free and Total

Test	Current Result and Flag	Previous Result and Date	Units	Reference Interval
Testosterone ⁰¹	387		ng/dL	264-916
Adult male reference interval is based on a population of healthy nonobese males (BMI <30) between 19 and 39 years old. Travison, et.al. JCEM 2017,102;1161-1173. PMID: 28324103.				
Free Testosterone(Direct) ⁰²	9.7		pg/mL	6.8-21.5

Estradiol

Test	Current Result and Flag	Previous Result and Date	Units	Reference Interval
Estradiol ⁰¹	19.5 Roche ECLIA methodology		pg/mL	7.6-42.6

Prostate-Specific Ag

Test	Current Result and Flag	Previous Result and Date	Units	Reference Interval
Prostate Specific Ag ⁰¹	0.7 Roche ECLIA methodology. According to the American Urological Association, Serum PSA should decrease and remain at undetectable levels after radical prostatectomy. The AUA defines biochemical recurrence as an initial PSA value 0.2 ng/mL or greater followed by a subsequent confirmatory PSA value 0.2 ng/mL or greater. Values obtained with different assay methods or kits cannot be used interchangeably. Results cannot be interpreted as absolute evidence of the presence or absence of malignant disease.		ng/mL	0.0-4.0

Disclaimer
The Previous Result is listed for the most recent test performed by Labcorp in the past 5 years where there is sufficient patient demographic data to match the result to the patient. Results from certain tests are excluded from the Previous Result display.

Icon Legend
▲ Out of Reference Range ■ Critical or Alert

Performing Labs

Patient Details	Physician Details	Specimen Details
Phone: Date of Birth: Age: Sex: Patient ID: Alternate Patient ID:	Request A Test, LTD. 7027 Mill Road Suite 201, BRECKSVILLE, OH, 44141 Phone: Physician ID: NPI:	Specimen ID: Control ID: Alternate Control Number: Date Collected: Date Received: Date Entered: Date Reported: Rte: